

Note to readers with disabilities: *EHP* strives to ensure that all journal content is accessible to all readers. However, some figures and Supplemental Material published in *EHP* articles may not conform to [508 standards](#) due to the complexity of the information being presented. If you need assistance accessing journal content, please contact ehp508@niehs.nih.gov. Our staff will work with you to assess and meet your accessibility needs within 3 working days.

Supplemental Material

Low Greenhouse Gas Emission Self-Selective Diets and Risk of Metabolic Syndrome in Adults 40 and Older: A Prospective Cohort Study in South Korea

Li-Juan Tan, and Sangah Shin

Table of Contents

Table S1. GHG emissions associated with Korean representative foods according to HEXA 106-item FFQ data.

Table S2. GHG emissions (kg CO₂ eq) per main food group in Korean Food Balance Wheel according to quintiles of energy-adjusted GHG emissions (N=41,659 from the HEXA Study, South Korea).

Table S3. Serving sizes (servings/day) per main food group in Korean Food Balance Wheel according to quintiles of energy-adjusted GHG emissions (N=41,659 participants in the HEXA Study, South Korea).

Table S4. MetS biomarkers collected at follow-up of the study participants from the Health Examinees Study according to the quintiles of the energy-adjusted diet-related GHG emissions ($n = 41,659$).

Figure S1. Flowchart of the selection of study participants.

Figure S2. Percentage proportions of GHG emissions per main food group in Korean Food Balance Wheel according to quintiles of energy-adjusted GHG emissions.

Figure S3. Serving proportions of the main food groups in Korean Food Balance Wheel according to quintiles of energy-adjusted GHG emissions.